

The occasional abdominal paracentesis

Peter Hutten-Czapowski,
MD

Professor of Family
Medicine, Northern Ontario
School of Medicine, Ontario,
Canada

Correspondence to:
Peter Hutten-Czapowski,
phc@srpc.ca

This article has been peer
reviewed.

INTRODUCTION

Once in a while, there is fluid in the abdomen. Usually, it is ascites from cirrhosis, heart failure or malignancy; it could even be infected. Which is it? (Imaging won't tell you.) A tap could make the diagnosis clear.¹ Can you make your patient more comfortable by draining most of it? Of course you can.²

EQUIPMENT

- Portable Ultrasound with abdominal probe and gel
- Surgical skin prep such as Provioline
- 5cc syringe and needles for anaesthetising the tract
- Xylocaine 1%
- Sterile drapes and gown
- #14-gauge IV cannula
- Three-way stop cock
- Sample tubes (typically for cells chemistry and cytology)
- Suction tubing and body of 10cc syringe as an adapter
- Wall suction with drainage carousel or evacuated vacuum containers

PROCEDURE

I was trained to do the procedure without imaging.

1. Position the patient on their side.

I go lateral to the rectus on the dependent side to avoid injury to the hypogastric artery. Avoid any scarred areas that might have adherent bowel

2. Mark the level of shifting dullness and go dependent to there. These days, I use an ultrasound, as a quick application of the probe will find a deep pocket of ascites for you clear of any vulnerable structure [Figure 1]. I'm old school, so after finding my spot, I mark it on the patient in pen. When doing the procedure, I can watch my hands and the patient, and not the screen. Some practitioners are agile enough to do the job two-handed, when watching the screen, with a sterile ultrasound probe cover (e.g., a sterile glove palm) in one hand and the needle in the other. Use the technique that you are comfortable with
3. Prep a generous area around the mark and drape the patient with sterile drapes
4. Using a small needle raise a skin bleb with 0.5 ml xylocaine [Figure 2]
5. Attach the syringe containing residual 5–10 ml xylocaine to a three-way stop cock and the back end of a 14-gauge intravenous (IV) cannula, infiltrating as you go [Figure 3]

Access this article online

Quick Response Code:



Website:
www.cjrm.ca

DOI:
10.4103/cjrm.cjrm_61_22

Received: 22-07-2022

Revised: 10-08-2022

Accepted: 11-08-2022

Published: 02-01-2023

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Hutten-Czapowski P. The occasional abdominal paracentesis. Can J Rural Med 2023;28:34-6.

6. Advance the needle slowly into the abdominal cavity alternately injecting local anaesthetic and applying slight suction. I place my finger along the needle to control the insertion [Figure 4]
7. When ascites fluid is aspirated, withdraw the steel needle and syringe from the cannula and replace the stopcock and syringe to draw samples (if desired). The arm of the stopcock always points to the port that is turned off. Note the colour and turbidity of the liquid. If turbid, infection or malignancy may be suspect. If diagnosis is the aim, samples can be drawn for cell count (purple top tube), protein and other chemistries (red top tube), cytology (formalin) and microbiology (usually a sterile urine container)
8. Attach the suction tubing to the unused port on the stopcock. Standard wall suction tubing

can be attached to a Luer lock connector with the body of a 10cc syringe [Figure 5]

9. Attach to wall suction and set on regulated low-to-medium suction [Figure 6]. The catheter can be easily displaced when waiting for the fluid to drain. Be careful not to withdraw all the fluids as this might drop the blood pressure.

COMPLICATIONS

Over a few decades of rural medical practice, I have encountered common complications of blocked needle, kinked needle, needle that gets pulled out and similar incidents that require replacement of the catheter. Occasionally, I have had a persistent fluid leak that may require a purse string closure and pressure dressing. I have never had a patient go hypotensive or bleed, but



Figure 1: Ultrasound showing large pocket of fluid in black under the probe



Figure 2: Freezing the skin



Figure 3: The cannula stopcock syringe assembly



Figure 4: Tapping




Figure 5: Capped adapter

if that occurs, I would treat with IV saline and potentially albumin. Bleeding from the abdominal wall can be controlled with a figure-of-eight suture. Intra-abdominal bleeding would require laparotomy.

CONCLUSION

Abdominocentesis can be easily done by the rural generalist physician to provide relief or diagnosis for your patient.

Acknowledgments: I thank Dr. Marie Lynn Lacasse who helped with the photography.



Figure 6: Suction collection

Financial support and sponsorship: Nil.

Conflicts of interest: There are no conflicts of interest.

REFERENCES

1. Romney R, Mathurin P, Ganne-Carrié N, Halimi C, Medini A, Lemaitre P, *et al.* Usefulness of routine analysis of ascitic fluid at the time of therapeutic paracentesis in asymptomatic outpatients. Results of a multicenter prospective study. *Gastroenterol Clin Biol* 2005;29:275-9.
2. Wittmer VL, Lima RT, Maia MC, Duarte H, Paro FM. Respiratory and symptomatic impact of ascites relief by paracentesis in patients with hepatic cirrhosis. *Arq Gastroenterol* 2020;57:64-8.

CALL FOR PAPERS

The Canadian Journal of Rural Medicine (CJRM) is a quarterly peer-reviewed journal available in print form and open access online. It is the first rural medical journal in the world indexed in Index Medicus, as well as MEDLINE/PubMed databases.

The CJRM seeks to promote research into rural health issues and promote the health of rural and remote communities. It looks to support and inform rural practitioners, provide a forum for debate and discussion of rural medicine, provide practical clinical information to rural practitioners and influence rural health policy by publishing articles that inform decision-makers.

The material in the following categories will be considered for publication.

- Original articles: research studies, case reports and literature reviews of rural medicine (3500 words or less, not including references)
- Commentary: editorials, regional reviews and opinion pieces (1500 words or less)
- Clinical articles: practical articles relevant to rural practice. Illustrations and photos are encouraged (2000 words or less)
- Off Call articles: a grab-bag of material of general interest to rural doctors (e.g., travel, musings on rural living, essays) (1500 words or less).
- Cover: artwork with a rural theme

For more information, please visit www.srpc.ca/cjrm