

Cai-Lei Matsumoto,
MSc¹, Sheldon Tobe,
MD, FRCP, MScCH^{2,7},
Yoko S. Schreiber, MD
FRCPC MSc (Epi)
CIP^{5,4}, Natalie Bock-
ing, MD, CCFP, RCP-
SC⁵, Janet Gordon,
PND¹, Sharen Madden,
MD, MSc, CCFP,
FCFP⁶, Josh Hopko¹,
Len Kelly, MD, MCl
Sci, FCFP, FRRM⁸

¹Sioux Lookout First Nations Health Authority,
²Division of Nephrology,
U of T Associate Scientist,
Institute of Medical Science,
University of Toronto,
³Division of Clinical Sciences,
Northern Ontario School of
Medicine at Sioux Lookout
Meno Ya Win Health Centre,
⁴Section of infectious diseases,
Max Rady College of Medi-
cine, University of Manitoba,
Winnipeg, ⁵Public Health
and Preventative Medicine
Specialist, Sioux Lookout
First Nations Health
Authority, Sioux Lookout,
Ontario, ⁶Division of Clinical
Sciences, Northern Ontario
School of Medicine, Thunder
Bay, Canada, ⁷Institute of
Medical Science, University
of Toronto, Toronto, ⁸Sioux
Lookout Meno Ya Win
Health Centre, Ontario

Correspondence to:
Len Kelly,
lkelly@mcmaster.ca

This article has been peer
reviewed.

Access this article online

Quick Response Code:



Website:
www.cjrm.ca

DOI:
10.4103/CJRM.CJRM_99_19

Diabetes prevalence and demographics in 25 First Nations communities in northwest Ontario (2014–2017)

Abstract

Introduction: First Nations communities are known to have high rates of diabetes. The rural First Nations communities in northwest (NW) Ontario are particularly affected. Regional studies in 1985 and 1994 found a high prevalence of diabetes. More recently, they are estimated to have the highest prevalence in Ontario at 19%, double the provincial norm. The purpose of this study is to examine the epidemiology and prevalence of diabetes in the total population and cardiovascular comorbidities in the adult population of 25 First Nations communities in NW Ontario.

Methods: This retrospective diabetes prevalence study used primary care electronic medical record data for a 3-year period, 1 August 2014–31 July 2017. Diabetes prevalence was calculated for both the total and the adult (18+) populations and comorbid hypertension and dyslipidaemia were identified in adults.

Results: The age-adjusted diabetes prevalence for the total population was 15.1% versus a Canadian prevalence of 8.8%. The age-adjusted adult prevalence was 14.1%, double Canada's average of 7.1%. The average age of adults with diabetes was 52 years (± 14.9); 57% were female. Comorbid hypertension (58%) and dyslipidaemia (73%) were common. Metformin was the most commonly used medication (58%), followed by insulin/analogues (23%) and sulphonylureas (13%).

Conclusion: The diabetes prevalence in the First Nations population of NW Ontario is double Canada's norm. Addressing it will require addressing relevant social determinants of health, including poverty and food security.

Keywords: Diabetes, First Nations, prevalence

Résumé

Introduction : Les communautés des Premières nations sont reconnues pour leur taux élevé de diabète, particulièrement les communautés rurales des Premières nations du Nord-Ouest de l'Ontario. Des études régionales réalisées en 1985 et 1994 ont révélé une forte prévalence de diabète. Plus récemment, on a estimé que la prévalence dans ces communautés s'élevait à 19 %, la plus forte en Ontario et le double de la norme provinciale. Cette étude visait à examiner l'épidémiologie et la prévalence du diabète auprès de la population totale et les comorbidités

Received: 22-11-2019 Revised: 10-12-2019 Accepted: 19-07-2020 Published: ***

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Matsumoto C, Tobe S, Schreiber YS, Bocking N, Gordon J, Madden S, et al. Diabetes prevalence and demographics in 25 First Nations communities in northwest Ontario (2014–2017). Can J Rural Med 2020;25:139-44.

cardiovasculaires auprès de la population adulte de 25 communautés des Premières nations du Nord-Ouest de l'Ontario.

Méthodologie : Cette étude rétrospective visant à évaluer la prévalence du diabète a eu recours aux données sur 3 ans des dossiers médicaux électroniques des cliniques de première ligne, soit du 1^{er} août 2014 au 31 juillet 2017. La prévalence du diabète a été calculée dans les populations totale et d'adultes (18 ans et plus) et l'hypertension et la dyslipidémie ont été dépistées en concomitance chez les adultes.

Résultats : La prévalence du diabète ajustée en fonction de l'âge dans la population totale était de 15,1 % par rapport à la prévalence canadienne de 8,8 %. La prévalence ajustée en fonction de l'âge chez les adultes était de 14,1 %, soit le double de la prévalence canadienne de 7,1 %. L'âge moyen des adultes diabétiques était de 52 ($\pm 14,9$) ans; et 57 % des participants étaient de sexe féminin. L'hypertension (58 %) et la dyslipidémie (73 %) étaient courantes en concomitance. La metformine était le médicament le plus fréquemment utilisé (58 %), suivie de l'insuline/analogues (23 %) et des sulfonurées (13 %).

Conclusion : La prévalence du diabète dans les populations des Premières nations du Nord-Ouest de l'Ontario est le double de celle du Canada. Pour régler la situation, il faudra se pencher sur les déterminants sociaux de la santé pertinents tels que la pauvreté et l'insécurité alimentaire.

Mots-clés : Diabète, Premières nations, prévalence

INTRODUCTION

First Nations communities are known to have high rates of diabetes, particularly the rural First Nations communities in northwest (NW) Ontario. Regional studies in 1985 and 1994 found a high prevalence of diabetes and in a 2019 International Credential Evaluation Service report (ICES), they are estimated to have the highest prevalence in Ontario, at 19%, double the provincial norm of 8%.¹⁻³ These communities also have higher rates of lower limb amputations ($\times 7$) and advanced kidney disease ($\times 2$).⁴⁻⁶ An accurate description of the regional scope of diabetes is needed to design appropriate initiatives and establish a baseline from which ongoing changes in disease presence can be measured. The purpose of this study is to examine the epidemiology and prevalence of diabetes in the total and adult population and associated cardiovascular comorbidities in diabetic adults in 25 First Nations communities in NW Ontario.

METHODS

Study design

This retrospective diabetes prevalence study used primary care electronic medical record (EMR) data for a 3-year period, 1 August 2014–31 July 2017. Diabetes prevalence was calculated for both the total and the adult (18+) population. Comorbid hypertension and dyslipidaemia prevalence was calculated for the adult population.

Setting and participants

The setting is 25 remote First Nations communities in NW Ontario, with a total population of 24,493, including 16,170 adults (18+). Each community receives primary care services from a local nursing station, with regular visits from community physicians, supported by the Sioux Lookout First Nations Health Authority (SLFNHA). Patients were included who had at least 1 primary care visit, at either one of the 25 nursing stations, or in the Sioux Lookout clinic, who were being investigated or treated for diabetes. Patients under age 18 were not included in the analysis of comorbidities.

Sources of data

Since 2013, all primary care clinical visits, laboratory investigations and prescriptions have been recorded in the OSCAR EMR in use by the practices of 30 community physicians. Data extraction and analysis was performed by the SLFNHA information technology department (JH) and verified by its public health epidemiologist (CM). The SLMHC laboratory performs all regional laboratory testing, with results automatically entered into the OSCAR. Medication prescribing is done exclusively through OSCAR and is linked to regional pharmacies.

Extracted data included demographics, laboratory results and prescribed medications. A 3-year time frame was chosen as most patients

with chronic conditions have at least one primary care visit in 2 years.^{7,8} Data collection protocol approximated the validated Canadian Primary Care Sentinel Surveillance Network methods, but used a glycated haemoglobin level (A1c) of 6.5% rather than 7.0%.^{9,10} Administrative data (physician billing and hospital diagnoses) were not used, but the use of EMR prescription and laboratory data has been shown to render equivalent results.⁷

Variables

Demographic and laboratory data included age, gender, low-density lipoprotein-cholesterol (LDL-C) and A1c. Prescribed medications were identified by the World Health Organization Anatomic Therapeutic Classification system.¹¹ This coding identifies all glucose and lipid-lowering and antihypertensive medications.

Diabetes was defined as a recorded A1c ≥ 6.5 mmol/l or a prescription of a glucose-lowering medication; hypertension was identified by the prescription of an antihypertensive medication and dyslipidaemia by an elevated LDL-C ≥ 2.0 mmol/L or the prescription of a lipid-lowering medication.

Statistical methods

The prevalence of diabetes was adjusted to the 2016 Canadian census population. The total population prevalence provided a comparator to national values, but most of the analysis focused on the adult prevalence, allowing comparison with other First Nations and the estimation of comorbid hypertension and dyslipidaemia. Data were presented as mean and standard deviation for continuous variables and proportions for discrete variables.

Ethics

SLFNHA requested and approved the study. The Sioux Lookout Meno Ya Win Research Review and Ethics Committee (#16-15) and the Lakehead University Research Ethics Board (#161 15-61) approved the study.

RESULTS

The age-adjusted population prevalence was 15.1%, with an adult (18+) prevalence of 14.4%. A total

of 2960 patients were identified with diabetes, including 2888 adults. A1c testing occurred in 32% of the total adult population (5161/16,170). Females received A1c testing more often, 57% (2942/5161), and constituted 57% of the adult diabetic population (1657/2888).

Adult diabetic patients had a mean age of 52 years, 11 years older than the average adult age and had double the rate of comorbid hypertension (58%) and dyslipidaemia (73%) [Table 1 and Figure 1].

The average adult A1c was 8.6 mmol/l; 29.5% were $\leq 7.0\%$. The most common glucose-lowering medication was metformin, followed by insulin and sulphonylureas [Table 2].

The greatest increase in age group prevalence occurred in females aged 30–40 years, from 8.9% at age 30–34 years to 20.2% at 35–39 years [Figures 2 and 3]. There was a wide range in the adult prevalence among the 25 communities; crude rates varied from 9% to 32%.

DISCUSSION

Prevalence

The 15.1% overall age-adjusted prevalence of diabetes in these 25 First Nations communities was higher than the 2017 Canadian prevalence of 8.8%.¹² While higher than national values, the prevalence is lower than that in other First Nations studies; our adult-specific age-adjusted prevalence of 14.4% is higher than the Canadian adult prevalence of 7.1%, but is lower than the crude adult prevalence of 27% in the James Bay Cree and 35% in a 2016 Manitoba First Nations screening study (our crude adult prevalence was 18%).^{13–15} Methodological variations exist in these comparator studies; both relied primarily on A1c laboratory data. However, they used different cohorts: James Bay prevalence was estimated in the context of a proactive community-based diabetic programme, whereas a self-referred screening population was used in Manitoba.^{14,15} It is not clear how these differences affect prevalence estimates, but First Nations communities in NW Ontario appear to have a lower diabetes prevalence than other First Nations. Our assumption that the 3-year study period we used would identify most patients with diabetes seemed appropriate; Grevier's Canada-wide primary care EMR-based study found that diabetic patients




Figure 1: Diabetes prevalence by age in 25 First Nations communities in northwest Ontario 2014-2017 and Canada, 2013-2014

Table 1: Demographics of total adult population and diabetic adult population in 25 First Nations communities, northwest Ontario, 2014-2017

Characteristics	Total adult population, n (%)	Diabetic adult population, n (%)
n (%)	16,170 (100)	2888 (18)
Mean age, SD	41 (16.9)	52 (14.9)
Female	8048 (50)	1657 (57)
Comorbidities		
Hypertension	3935 (24)	1675 (58)
Dyslipidaemia	5210 (32)	2108 (73)

SD: Standard deviation

Table 2: Adult use of diabetes medications (n=2888)

Class of drug	%
Biguanides (metformin)	58
Insulin/analogues	23
Sulphonylureas	13
DPP-4 inhibitors	3
Other medications	2
No medication	7

DPP-4: Dipeptidyl peptidase 4

were 1.4 times more likely to have a primary care visit in their 2-year study period and the diabetes programme in James Bay Cree communities found that 90% of the patients with diabetes had an A1c measurement within a 2-year period.^{7,15}

Age and gender

First Nations populations are known to develop diabetes at an earlier age than other Canadians.¹⁶ The age-grouped results support this, demonstrating higher prevalence at younger ages [Figure 1]. Women were equally overrepresented (57%) in both the A1c-tested and




Figure 2: Adult (18+) diabetes prevalence by age and gender in 25 First Nations communities, northwest Ontario, 2014-2017.

diabetic populations. The largest diabetic cohort increase occurred in females in their 30s, where it doubled. Our regional population has a high birth rate, a high incidence of gestational diabetes and a high incidence of transition to overt Type 2 diabetes following gestational diabetes.¹⁷⁻¹⁹ These contribute to the increased female prevalence of diabetes after age 30 and are consistent with a 1985 study of the same communities and other recent First Nations studies, which demonstrated a female preponderance²⁰⁻²⁴ [Figure 2]. This contrasts with national values with a higher prevalence in males²⁵ [Figure 3].

Glycaemic control

The average A1c value in this adult diabetic population was 8.6%, higher than the 7.3%–8.2% in other primary care studies, with First Nations populations generally being at the higher end of that range^{15,26-28} The target glycaemic control of A1c $\leq 7.0\%$ was achieved in 29.5% of patients, compared to 39%–51% in other primary care studies.^{28,29} Similarly, metformin was the most commonly prescribed glucose-lowering medication.^{29,30}

Comorbidities

Hypertension co-prevalence (58%) in First Nations populations was similar to that in other Canadian diabetic populations (67.1%), but well below the 92% found in a 2011 study of 19 Canadian First Nations communities, whose methods included blood pressure measurement data in addition to antihypertensive medication records for 885 diabetic patients.^{26,31} Dyslipidaemia co-prevalence of 73% was




Figure 3: Diabetes prevalence (cases/100) by age and gender in 25 First Nations communities in northwest Ontario 2014–2017 and Canada, 2013–2014.

between the 59% in Canada and 93% in other First Nations studies.^{27,31}

Context

Estimates of diabetes prevalence tell only part of the story. The First Nations population experience increased diabetes-related morbidity and mortality.³² Many factors contribute to this disease burden, including cultural and social determinants of health, poverty and food insecurity. Historic and political factors include inequities in employment and education, marginalisation, racism and intergenerational trauma resulting from governmental assimilation practices.^{33–37} A similar disease profile exists in culturally and geographically distinct First Nations communities across Canada which reflects the impact of this shared history of colonisation and its ongoing detrimental effects.^{33,36,37} Addressing these factors requires broad social and healthcare responses. The wide range of community-specific prevalence requires further examination to understand how genetic, epigenetic or environmental factors affect rates of diabetes.³⁸

Limitations

A variety of methods and data are used in diabetes prevalence research. The Public Health Agency of Canada uses administrative data (hospital and physician diagnoses). We used laboratory and medication EMR data. Diagnostic codes for primary care visits were not available to us, as they are not automatically recorded in this EMR, which

also precluded distinguishing between different classes of diabetes. We assumed a stable population as most communities are geographically isolated; but participation in the study did include northern patients who have moved to Sioux Lookout and some Sioux Lookout area residents.

CONCLUSION

The 25 First Nations communities in NW Ontario have a diabetes prevalence twice the Canadian average. Patients diagnosed with diabetes were younger, included more females and had significant levels of comorbid hypertension and dyslipidaemia. Social determinants of health and food security will need a culturally appropriate broad-based approach to address the environment in which diabetes thrives.

Financial support and sponsorship: This work was supported by the Northern Ontario Academic Medicine Association.

Conflicts of interest: There are no conflicts of interest.

REFERENCES

1. Young TK, McIntyre LL, Dooley J, Rodriguez J. Epidemiologic features of diabetes mellitus among Indians in northwestern Ontario and northeastern Manitoba. *Can Med Assoc J* 1985;132:793–7.
2. Fox C, Harris SB, Whalen-Brough E. Diabetes among Native Canadians in Northwestern Ontario: 10 Years later. *Chronic Dis Canada* 1994;15:92–6.
3. Green ME, Jones CR, Walker JD, Shah BR, Jacklin K, Slater M, *et al.*, editors. *First Nations and Diabetes in Ontario*. Toronto, ON: ICES; 2019.
4. Tobe S, Gordon J, Hopko J, Madden S, Willms H, Kelly L. Chronic kidney disease prevalence in first nations communities

in NW Ontario: Prevalence of CKD and comorbidities. *CMAJ Open* 2019;7:E568-72.

5. Loewen K, Vigliarolo J, Lance B, Rockley M, Schreiber Y, Kivi C, *et al.* Diabetes, foot ulcers, amputation rates in NW Ontario: An incidence study and the introduction of a standardized diabetic foot ulcer management protocol. *CJRM* 2017;22:100-7.
6. Booth GL, Polsky JY, Gozdyra G, Cauch-Dudek K, Kiran T, Shah BR, *et al.* Regional Measures of Diabetes Burden in Ontario. Toronto: Institute for Clinical Evaluative Sciences; 2012. Available from: <https://www.ices.on.ca/Publications/Atlases-and-Reports/2012/Regional-Measures-of-Diabetes-Burden-in-Ontario>. [Last accessed on 2019 Nov 25].
7. Greiver M, Williamson T, Bennett TL, Drummond N, Savage C, Aliarzadeh B, *et al.* Developing a method to estimate practice denominators for a national Canadian electronic medical record database. *Fam Pract* 2013;30:347-54.
8. Sibley LM, Moineddin R, Agha MM, Glazier RH. Risk adjustment using administrative data-based and survey-derived methods for explaining physician utilization. *Med Care* 2010;48:175-82.
9. CPCSSN Disease Definitions Version 2. Kingston (ON): Canadian Primary Care Sentinel Surveillance Network; 2019. Available from: <http://cpcssn.ca/wp-content/uploads/2019/07/CPCSSN-Case-Definitions-v2.pdf>. [Last accessed on 2020 Aug 05]
10. Williamson T, Green ME, Birthwhistle R, Khan S, Garies S, Wong S, *et al.* Validating the 8 CPSSN case definitions for chronic disease in a primary care database of electronic health records. *Ann Fam Med* 2014;12:367-72.
11. Guidelines for ATC Classification and DDD Assignment 2019. Oslo: WHO Collaborating Centre for Drug Statistics Methodology; 2018. Available from: http://www.whocc.no/filearchive/publications/_guidelines_web.pdf. [Last accessed on 2019 Nov 27].
12. Public Health Agency of Canada. At a Glance- Twenty Years of Diabetes Surveillance Using the Canadian Chronic Surveillance System. Available from: <https://www.canada.ca/en/public-health/services/reports-publications/health-promotion-chronic-disease-prevention-canada-research-policy-practice/vol-39-no-11-2019/twenty-years-diabetes-surveillance.html>. [Last accessed on 2019 Nov 22].
13. Statistics Canada. Diabetes by age Group. Available from: <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310009607>. [Last accessed on 2019 Nov 20].
14. Komenda P, Lavallee B, Ferguson TW, Tangri N, Chartrand C, McLeod L, *et al.* The prevalence of CKD in rural Canadian indigenous peoples: Results from the First Nations community based screening to improve kidney health and prevent dialysis (FINISHED) Screen, triage, and treat program. *Am J Kidney Dis* 2016;68:582-90.
15. Cree Board of Health and Social Services of James Bay. Summary Health Profile March 11, 2016. Background Summary Statistical Profile of Health and Health Determinants in Eeyou Istchee for the Health Assembly. Available from: <https://www.creehealth.org/sites/default/files/A%20Health%20Profile%20for%20Health%20Assembly.pdf>. [Last accessed on 2019 Nov 05].
16. Oster RT, Johnson JA, Balko SU, Svenson LW, Toth EL. Increasing rates of diabetes amongst status Aboriginal youth in Alberta, Canada. *Int J Circumpolar Health* 2012;71:1-7.
17. Hummelen R, Kattini R, Poirier J, Kelly L, Madden S, Ockenden H, Dooley J. Demographics, prevalence and outcomes of diabetes in pregnancy in NW Ontario and small Ontario obstetrical programs. *CJRM* 2020;3:99-104.
18. Sioux Lookout First Nations Health Authority. Our Children and Youth Health Report. Sioux Lookout, Ontario; 2018. Available from: https://slfnha.com/application/files/62154022/0883/CHSR_FINAL_-_Web_Version.pdf. [Last accessed on 2019 Jun 10].
19. Mohamed N, Dooley J. Gestational diabetes and subsequent development of NIDDM in aboriginal women of northwestern Ontario. *Int J Circumpolar Health* 1998;57 Suppl 1:355-8.
20. Hemond J, Robbins RB, Young PC. The Effects of Maternal Obesity on Neonates, Infants, Children, Adolescents, and Adults. *Clin Obstet Gynecol* 2016;59:216-27.
21. Harris S, Naqshbandi M, Bhattacharyya O, Hanley A, Esler J, Zinman B On behalf of the CIRCLE Study Group Major gaps in diabetes clinical care among Canada's First Nations: Results of the CIRCLE study. *Diabetes Res Clin Practice* 2011;92:272-9.
22. Dyck R, Osgood N, Lin TH, Gao A, Stang MR. Epidemiology of diabetes mellitus among First Nations and non-First Nations adults. *CMAJ* 2010;182:249-56.
23. Oster RT, Johnson JA, Hemmelgarn BR, King M, Balko SU, Svenson LW, *et al.* Recent epidemiologic trends of diabetes mellitus among status Aboriginal adults. *CMAJ* 2011;183:E803-8.
24. Young TK, Reading J, Elias B, O'Neil JD. Type 2 diabetes mellitus in Canada's First Nations: Status of an epidemic in progress. *CMAJ* 2000;163:561-6.
25. Statistics Canada. Health Fact Sheet. Diabetes; 2017. Available from: <https://www150.statcan.gc.ca/n1/pub/82-625-x/2018001/article/54982-eng.htm>. [Last accessed on 2019 Nov 19].
26. Harris S, Naqshbandi MM, Bhattacharyya O. Major gaps in diabetes clinical care among Canada's First Nations: Results of the CIRCLE study. *Diabetes Res Clin Pract* 2011;92:272-9.
27. Harris SB, Kapor J, Lank CN, Willan AR, Houston T. Clinical inertia in patients with T2DM requiring insulin in family practice. *Can Fam Physician* 2010;56:e418-24.
28. Leiter L, Berard L, Bowering K, Cheng Y, Dawson K, Ekoe J, *et al.* Type 2 diabetes mellitus in Canada: Is it improving? *Canadian J Diabetes* 2013;37:82-9.
29. Greiver M, Williamson T, Barber D, Birtwhistle R, Aliarzadeh B, Khan S, *et al.* Prevalence and epidemiology of diabetes in Canadian primary care practices: A report from the Canadian Primary Care Sentinel Surveillance Network. *Can J Diabetes* 2014;38:179-85.
30. Spann SJ, Nutting PA, Galliher JM, Peterson KA, Pavlik VN, Dickinson LM, *et al.* Management of type 2 diabetes in the primary care setting: A practice-based research network study. *Ann Fam Med* 2006;4:23-31.
31. Padwal R, Bienek A, McAlister F, Campbell N. Research task force of the Canadian hypertension education program. Epidemiology of hypertension in Canada: An update. *Canadian J Cardiol* 2016;32:687-94.
32. Crowshoe L, Dannenbaum D, Green M, Henderson R, Naqshbandi Hayward M, *et al.* Diabetes Canada clinical practice guidelines expert committee. Diabetes Canada 2018 clinical practice guidelines for the prevention and management of diabetes in Canada" Type 2 diabetes and indigenous peoples. *Can J Diabetes* 2018;42 Suppl 1:S1-325.
33. Allan B, Smylie J. First Peoples, Second Class Treatment: The role of Racism in the Health and Wellbeing of Indigenous Peoples in Canada. Toronto: Wellesley Institute; 2015.
34. An overview of Aboriginal Health. National Collaborating Centre for Indigenous Health 2013. Prince George B.C. Available from: <https://www.ccnsa-nccah.ca/docs/context/FS-OverviewAboriginalHealth-EN.pdf>. [Last accessed on 2019 Aug 08].
35. Jacklin KM, Henderson RI, Green ME, Walker LM, Calam B, Crowshoe LJ. Health care experiences of Indigenous people living with type 2 diabetes in Canada. *CMAJ* 2017;189:E106-12.
36. Halseth R. The Prevalence of Type 2 Diabetes among First Nations and Considerations for Prevention. Prince George, BC: National Collaborating Centre for Aboriginal Health; 2019.
37. Smylie J, Phillips-Beck W. Truth, respect and recognition: Addressing barriers to Indigenous maternity care. *CMAJ* 2019;191:E207-E208.
38. Sterns JD, Smith CB, Steele JR, Stevenson KL, Gallicano GI. Epigenetics and type II diabetes mellitus: Underlying mechanisms of prenatal predisposition. *Front Cell Dev Biol* 2014;2:15.