

The occasional C-spine x-ray

Karl Stobbe, MD, CCFP (EM)
Beamsville, Ont.

CJRM 2004;9(1):38-42

Please turn to [page 42](#) for the C-spine X-ray Checklist.

Introduction

X-rays are used in conjunction with the clinical examination to "clear" the c-spine in patients who present with multiple trauma, head injuries with neck pain, and those who have altered levels of consciousness and unknown injuries. Recently, the "Canadian C-spine Rule" was published to help identify patients who may NOT require a c-spine x-ray.¹

A lateral c-spine view can detect 60% to 80% of fractures.² Anterior-posterior (A-P) and open-mouth odontoid views improve the sensitivity to 80% to 95% or better.^{3,4} For those of us who only occasionally interpret c-spine films, oblique views can make it much easier to detect certain injuries. Occasionally, fractures poorly seen on other views will be seen with flexion or extension. Adding a CT scan increases the sensitivity to 95% to 100%;^{2,3} however, plain films are better than CT for detecting some injuries.⁵

Serious multiple injuries often require urgent treatment. Frequently, the c-spine is not cleared until after the patient is transferred to the referral centre.

Classification of C-spine fractures

For those who infrequently read these x-rays, there are 2 types of fractures: easy to find and hard to find.

1. Easy to find:

- C3-C6 flexion/extension injuries.

2. Hard to find:

- rotatory subluxations
- upper c-spine fractures - C1 & C2
- fractures in non-visualized C6 or C7

Most of these injuries can be detected or suspected in a lateral c-spine film and confirmed on other views.

Systematic interpretation

A systematic approach can help the physician to confidently determine whether a radiographic abnormality is visible.

Step A

Start with lateral c-spine ([Fig. 1](#)). Leave the patient on the stretcher or backboard with the hard collar on while this x-ray is being taken.

A1

Check for adequacy of the film. All of C7, and the top of T1 should be visible. The lower levels need to be visualized before the patient can be discharged with a cleared c-spine.⁶ On the other hand, if a fracture is visible on an "inadequate" film there is no need to risk further injury by moving the patient to get more views.

A2

Check the alignment. Follow the 4 lines indicated on the diagram. Line 1 is formed from the anterior aspect of the vertebral bodies and continues to the top of the odontoid process. The anterior arch of C1 lies anterior to this line. Osteophytes often create irregularities in this line. Line 2 follows the posterior aspect of the vertebral bodies and meets Line 1 at the top of the odontoid. This line is not usually affected by osteophytes. Line 3 connects the laminae, and also marks the posterior border of the spinal canal. Line 4 connects the tips of the spinous processes. Note that C1 does not touch this line.

A3

Look for soft tissue swelling ([Fig. 2](#)). In adults, the upper limit of normal is:

- 5 mm at C2
- 22 mm at C6

Next, check the space between the odontoid and C1 (atlas) anterior to it. This "atlanto-odontoid" space should be 3 mm or less.

A4

Look at the bones and disk-spaces. From C3-C7 each vertebral body should be about the same shape and size as its neighbour above and below. If the odontoid is intact on this film and the 4 lines demonstrate no abnormality, it is reasonable to proceed to more views. If normal but limited by not seeing C7, an attendant standing at the foot of the bed can pull on the patient's hands to bring the shoulders down.

If C7 cannot be seen despite arm traction, a swimmer's view will be required. To find C6 and C7 on this view, look for vertebrae articulating with ribs, indicating thoracic levels. C6 and C7 can be identified by counting up from T1.

Step B

If the lateral is normal; remove immobilization and proceed with more views.

A-P ([Fig. 3](#)):

- look for alignment of the spinous processes

Odontoid view ([Fig. 4](#)):

- Check alignment of the lateral masses
- Look for odontoid fracture

Step C

Obliques ([Fig. 5](#)) are not routinely done in all hospitals. These can be useful to show unilateral facet dislocations. Look for:

- Intact foraminae. Facet dislocations often obliterate one foramen on this view.
- The laminae should line up like "shingles on a roof." This alignment is disrupted in a facet dislocation.

Flexion and extension views occasionally show injuries not otherwise seen.⁷

Interpretation follows the same system as for the initial lateral c-spine view. These views can be falsely reassuring when the patient cannot adequately flex or extend the neck.⁸

Rural physicians are regularly faced with patients who have suffered trauma and have neck pain. Plain x-rays are usually the only diagnostic test available in the rural hospital. Plain films of the cervical spine have good sensitivity for fractures when compared with CT scanning. A systematic approach to interpretation can improve the confidence of rural physicians in interpreting these films. This systematic approach is summarized in [Figure 6](#), the Checklist on page 42.

Rural Program Coordinator, Department of Family Medicine, McMaster University, Hamilton, Ont.

Competing interests: None declared.

This article has been peer reviewed.

Correspondence to: Dr. Karl Stobbe, Box 430, Beamsville ON L0R 1B0

References

1. Stiell IG, Wells GA, McKnight RD, Brison R, Lesiuk H, Clement CM, et al. Canadian C-Spine Rule study for alert and stable trauma patients: I. Background and rationale. CJEM 2002;4(2):84-90. Available: www.caep.ca/004.cjem-jcmu/004-00.cjem/vol-4.2002/v42-084.htm#main (accessed 2004 Jan 5).
2. Crim JR, Moore K, Bordke D. Clearance of the cervical spine in multitrauma patients: the role of advanced imaging. Semin Ultrasound CT MR 2001;22(4):283-305.
3. Borock EC, Gabram SG, Jacobs LM, Murphy MA. A prospective analysis of a two-year experience using computed tomography as an adjunct for cervical spine clearance. J Trauma 1991;31(7):1001-5; discussion 1005-6.
4. Mace SE. Emergency evaluation of cervical spine injuries: CT versus plain radiographs. Ann Emerg Med 1985;14(10):973-5.
5. Woodring JH, Lee C. The role and limitations of computed tomographic scanning in the evaluation of cervical trauma. J Trauma 1992;33(5):698-708.
6. Davis JW, Phreaner DL, Hoyt DB, Mackersie RC. The etiology of missed cervical spine injuries. J Trauma 1993;34(3):342-6.
7. Lewis LM, Docherty M, Ruoff BE, Fortney JP, Keltner RA Jr, Britton P. Flexion-extension views in the evaluation of cervical-spine injuries. Ann

Emerg Med 1991;20(2):117-21.

8. Insko EK, Gracias VH, Gupta R, Goettler CE, Gaieski DF, Dalinka MK. Utility of flexion and extension radiographs of the cervical spine in the acute evaluation of blunt trauma. J Trauma 2002;53(3):426-9.

© 2004 Society of Rural Physicians of Canada




C-spine x-ray checklist (adults)

A. Lateral c-spine – keep hard collar on.

- ☐ adequacy of film – see all of C7, top of T1. If “No” – continue anyway. Repeat films if no fracture.
- ☐ Alignment – line 1 – continues along ant. odontoid
- ☐ Line 2 – continues along posterior odontoid
- ☐ Line 3
- ☐ Line 4 note C1 should *not* reach this line
- ☐ Soft tissue thickness < 5 mm at C2
- ☐ < 22 mm at C6
- ☐ atlanto–odontoid space: 3 mm or less
- ☐ Bones – check for fractures. Shape of C3–C7 vertebral body should be approx. the same as the one above and below.


If a fracture is seen – do no further views – immobilize and transfer

B. If lateral is normal, remove immobilization and proceed with more views



Odontoid view:


- ☐ Alignment of lateral masses (arrows)
- ☐ Odontoid fracture – 3 types shown



PA:

- ☐ Alignment of spinous processes only

- ### C. Obliques – Useful for unilateral facet dislocations
- ☐ Foraminae all intact (F)
 - ☐ Laminae line up like “shingles on a roof”



D. **Flexion–extension views** – for further reassurance if other views normal. Use lat c-spine checklist. Some injuries not obvious on initial lateral will be visible on a lateral view.

Fig. 6. C-spine x-ray checklist (adults)

[\[Return to text\]](#)



Fig. 1. Lateral c-spine view

[\[Return to text\]](#)




Fig. 2. Lateral c-spine:

- Adequate film: all of C7 is visible, as well as the top of T1
- All 4 lines form smooth arcs. Note that Lines 1 and 2 meet at the top of the odontoid (O). Note also that posteriorly C1 does not touch Line 4, and C1 (A) lies anterior to Line 1.
- Atlanto-odontoid space between C1 (A) and odontoid (O) is 3 mm or less.
- Soft tissue thickness anterior to C2 (B) is 5 mm or less.
- Soft tissue thickness anterior to C6 (C) is 22 mm or less.

[\[Return to text\]](#)



Fig. 3. A-P view: Note the alignment of spinous processes.

[\[Return to text\]](#)




Fig. 4. Odontoid view: the lateral masses of C1 line up with the corresponding structure on C2 (indicated by arrows).

[\[Return to text\]](#)




Fig. 5. Oblique view: Note the open neural foraminae (F), and the laminae (arrow) lining up like “shingles on a roof.”

[\[Return to text\]](#)


C-spine x-ray checklist (adults)

A. Lateral c-spine – keep hard collar on.

- ☐ adequacy of film – see all of C7, top of T1. If “No” – continue anyway. Repeat films if no fracture.
- ☐ Alignment – line 1 – continues along ant. odontoid
- ☐ Line 2 – continues along posterior odontoid
- ☐ Line 3
- ☐ Line 4 note C1 should *not* reach this line
- ☐ Soft tissue thickness < 5 mm at C2
- ☐ < 22 mm at C6
- ☐ atlanto–odontoid space: 3 mm or less
- ☐ Bones – check for fractures. Shape of C3–C7 vertebral body should be approx. the same as the one above and below.


If a fracture is seen – do no further views – immobilize and transfer

B. If lateral is normal, remove immobilization and proceed with more views



Odontoid view:


- ☐ Alignment of lateral masses (arrows)
- ☐ Odontoid fracture – 3 types shown



PA:

- ☐ Alignment of spinous processes only

- ### C. Obliques – Useful for unilateral facet dislocations
- ☐ Foraminae all intact (F)
 - ☐ Laminae line up like “shingles on a roof”



D. Flexion–extension views – for further reassurance if other views normal. Use lat c-spine checklist. Some injuries not obvious on initial lateral will be visible on a lateral view.

Fig. 6. C-spine x-ray checklist (adults)

[\[Return to text\]](#)

Country cardiograms case 24: Interpretation and discussion

Charles Helm, MD
Tumbler Ridge, BC

CJRM 2004;9(1):48-9

The first ECG (Fig. 1) shows sinus bradycardia, rate 54 beats/min. All intervals are normal, as is the QRS axis. The only area of possible concern lies in leads VI and V2. In V1, the magnitude of the height of the R wave is approximately equal to that of the depth of the S wave, and in V2 a much taller R wave is present.

Three common possibilities should be considered when this type of phenomenon is encountered: right ventricular hypertrophy, posterior myocardial infarction or counter-clockwise rotation.

Right ventricular hypertrophy is excluded in this case by the absence of any other supporting ECG features.

In many cases it is not possible to distinguish between counterclockwise rotation and isolated old posterior myocardial infarction. In this case, in the absence of an old tracing for comparison, we cannot tell whether the prominent R waves are old or new. However, ST-T segment changes, if present, would point to an ischemic process. The ST segment in V1 in the first ECG (Fig. 1), shows possibly some slight ST depression and T-wave inversion. Yet T-wave inversion is normal in this lead, and the possible ST-segment depression is probably minor enough not to cause undue concern.